## **AP CALCULUS PROBLEM SET #5 Motion (s-v-a)**

(97-1)

- 1. A particle moves along the x-axis so that its velocity at any time  $t \ge 0$  is given by  $v(t) = 3t^2 2t 1$ . The position x(t) is 5 for t = 2.
- (a) Write a polynomial expression for the position of the particle at any time  $t \ge 0$ .
- (b) For what values of t,  $0 \le t \le 3$ , is the particle's instantaneous velocity the same as its average velocity on the closed interval [0, 3]?
- (c) Find total distance travelled by the particle from time t = 0 until time t = 3.

(93-2)


- 2. A particle moves on the x-axis so that its position at any time t > 0 is given by  $x(t) = 2te^{-t}$ .
- (a) Find the acceleration of the particle at t = 0.
- (b) Find the velocity of the particle when its acceleration is 0.
- (c) Find the total distance travelled by the particle from t = 0 to t = 5.

(2012-6)

- 3. For  $0 \le t \le 12$ , a particle moves along the *x*-axis. The velocity of the particle at time *t* is given by  $v(t) = \cos\left(\frac{\pi t}{6}\right)$ . The particle is at position x = -2 at time t = 0.
- (a) For  $0 \le t \le 12$ , when is the particle moving to the left?
- (b) Write, but do not evaluate, an integral expression that gives the total distance traveled by the particle from time t = 0 to time t = 6.
- (c) Find the acceleration of the particle at time t. Is the speed of the particle increasing, decreasing, or neither at time t = 4? Explain your reasoning.
- (d) Find the position of the particle at time t = 4.

(2010(B)-6)

- 4. Two particles move along the x-axis. For  $0 \le t \le 6$ , the position of the particle P at time t is given by  $p(t) = 2\cos\left(\frac{\pi}{4}t\right)$ , while the position of particle R at time t is given by  $r(t) = t^3 6t^2 + 9t + 3$ .
- (a) For  $0 \le t \le 6$ , find all times t during which the particle R is moving to the right.
- (b) For  $0 \le t \le 6$ , find all times t during which the two particles travel in opposite directions.
- (c) Find the acceleration of particle P at time t = 3. Is particle P speeding up, slowing down, or doing neither at time t = 3? Explain your reasoning.
- (d) Write, but do not evaluate, an expression for the average distance between the two particles on the interval  $1 \le t \le 3$ .




Graph of v

A particle moves along the x-axis so that its velocity at time t, for  $0 \le t \le 6$ , is given by a differentiable function v whose graph is shown above. The velocity is 0 at t = 0, t = 3, and t = 5, and the graph has horizontal tangents at t = 1 and t = 4. The areas of the regions bounded by the t-axis and the graph of v on the intervals [0,3], [3,5], and [5,6] are 8, 3, and 2, respectively. At time t = 0, the particle is at x = -2.

- (a) For  $0 \le t \le 6$ , find both the time and the position of the particle when the particle is farthest to the left. Justify your answer.
- (b) For how many values of t, where  $0 \le t \le 6$ , is the particle at x = -8? Explain your reasoning.
- (c) On the interval  $2 \le t \le 3$ , is the speed of the particle increasing or decreasing? Give a reason for your answer.
- (d) During what time intervals, if any, is the acceleration of the particle negative? Justify your answer.

(2000-2) **6.** 




Two runners, A and B, run on a straight racetrack for  $0 \le t \le 10$  seconds. The graph above, which consists of two line segments, shows the velocity, in meters per second, of Runner A. The velocity, in meters per second, of Runner B is given by the function v defined by  $v(t) = \frac{24 t}{2t+3}$ .

- (a) Find the velocity of Runner A and the velocity of Runner B at time t=2 seconds. Indicate units of measure.
- (b) Find the acceleration of Runner A and the acceleration of Runner B at time t=2 seconds. Indicate units of measure.
- (c) Find the total distance run by Runner A and the total distance run by Runner B over the time interval  $0 \le t \le 10$  seconds. Indicate units of measure.

## **OPTIONAL**

(2001-3) 7.



A car is traveling on a straight road with velocity 55 ft/sec at time t = 0. For  $0 \le t \le 18$  seconds, the car's acceleration a(t), in ft/sec<sup>2</sup>, is the piecewise linear function defined by the graph above.

- (a) Is the velocity of the car increasing at t = 2 seconds? Why or why not?
- (b) At what time in the interval  $0 \le t \le 18$ , other than t = 0, is the velocity of the car 55 ft/sec? Why?
- (c) On the time interval  $0 \le t \le 18$ , what is the car's absolute maximum velocity, in ft/sec, and at what time does it occur? Justify your answer.
- (d) At what times in the interval  $0 \le t \le 18$ , if any, is the car's velocity equal to zero? Justify your answer.

(99-1)

- 8. A particle moves along the y-axis with velocity given by  $v(t) = t \sin(t^2)$  for  $t \ge 0$ .
- (a) In which direction (up or down) is the particle moving at time t = 1.5? Why?
- (b) Find the acceleration of the particle at time t = 1.5. Is the velocity of the particle increasing at t = 1.5? Why or why not?
- (c) Given that y(t) is the position of the particle at time t and that y(0) = 3, find y(2).
- (d) Find the total distance traveled by the particle from t = 0 to t = 2.

(76BC-1)

- 9. A particle moves on the x-axis in such a way that its position at time t is given by  $x = (2t-1)(t-1)^2.$
- (a) At what times t is the particle at rest?
- (b) During what interval of time is the particle moving to the left? Justify your answer.
- (c) At what time during the interval found in (b) is the particle moving most rapidly (that is, the speed is a maximum)? Justify your answer.