AP CALCULUS PROBLEM SET #8 CURVE ANALYSIS II

(77-2)


- Consider the function f defined by $f(x) = (x^2 1)^3$ for all real numbers x. 1.
- (a) For what values of x is the function increasing?
- (b) Find the x- and y-coordinates of the relative maximum and minimum points. Justify your answer.
- (c) For what values of x is the graph of f concave upward?
- (d) Using the information found in parts (a), (b), and (c), sketch the graph of f on the axes provided.

(89BC-3)

- Consider the function f defined by $f(x) = e^x \cos x$ with domain $[0, 2\pi]$.
- (a) Find the absolute maximum and minimum values of f(x).
- (b) Find the intervals on which f is increasing.
- (c) Find the x-coordinate of each point of inflection of the graph of f.

(2015-5)

3.



The figure above shows the graph of f', the derivative of a twice-differentiable function f, on the interval [-3, 4]. The graph of f' has horizontal tangents at x = -1, x = 1, and x = 3. The areas of the regions bounded by the x-axis and the graph of f' on the intervals [-2, 1] and [1, 4] are 9 and 12 respectively.

- (a) Find all x-coordinates at which f has a relative maximum. Give a reason for your answer.
- (b) On what open intervals contained in -3 < x < 4 is the graph of f both concave down and decreasing? Give a reason for your answer.
- (c) Find the x-coordinates of all points of inflection for the graph of f. Give a reason for your answer.
- (d) Given that f(1) = 3, write an expression for f(x) that involves an integral. Find f(4) and f(-2).

- 4. Let f be the function given by $f(x) = \frac{\ln x}{x}$ for all x > 0. The derivative of f is given by $f'(x) = \frac{1 \ln x}{x^2}$.
- a) Write an equation for the line tangent to the graph of f at $x = e^2$.
- b) Find the x-coordinate of the critical point of f. Determine whether this point is a relative minimum, a relative maximum or neither for the function f. Justify your answer.
- c) The graph of the function f has exactly one point of inflection. Find the x-coordinate of this point.
- d) Find $\lim_{x\to 0^+} f(x)$

(92BC-4)

5. Let
$$f$$
 be a function defined by $f(x) = \begin{cases} 2x - x^2 & \text{for } x \le 1 \\ x^2 + kx + p & \text{for } x > 1 \end{cases}$


- (a) For what values of k and p will f be continuous and differentiable at x = 1?
- (b) For the values of k and p found in part (a), on what interval or intervals is f increasing?
- (c) Using the values of k and p found in part (a), find all points of inflection of the graph of f. Support your conclusion.

(2001-4)

- 6. Let h be a function defined for all $x \ne 0$ such that h(4) = -3 and the derivative h is given by $h'(x) = \frac{x^2 2}{x}$ for all $x \ne 0$.
- (a) Find all values of x for which the graph of h has a horizontal tangent, and determine whether h has a local maximum, a local minimum, or neither at each of these values. Justify your answers.
- (b) On what intervals, if any, is the graph of h concave up? Justify your answer.
- (c) Write an equation for the line tangent to the graph of h at x = 4.
- (d) Does the line tangent to the graph of h at x = 4 lie above or below the graph of h for x > 4? Why?

OPTIONAL

(2013-4) **7.**



The figure above shows the graph of f', the derivative of a twice-differentiable function f, on the closed interval $0 \le x \le 8$. The graph of f' has horizontal tangent lines at x = 1, x = 3, and x = 5. The areas of the regions between the graph of f' and the x-axis are labeled in the figure. The function f is defined for all real numbers and satisfies f(8) = 4.

- (a) Find all values of x on the open interval $0 \le x \le 8$ for which the function f has a local minimum. Justify your answer.
- (b) Determine the absolute minimum value of f on the closed interval $0 \le x \le 8$. Justify your answer.
- (c) On what open intervals contained in $0 \le x \le 8$ is the graph of f both concave down and increasing? Explain your reasoning.
- (d) The function g is defined by $g(x) (f(x))^3$. If $f(3) = -\frac{5}{2}$, find the slope of the line tangent to the graph of g at x = 3.